

THE UNITED STRATES OF ANTERRICA

TO ALL TO WHOM THESE PRESENTS SHALL COME;

Pioneer Hi-Bred International, Inc.

MICCORS, THERE HAS BEEN PRESENTED TO THE

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE RIGHT TO EXCLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR IMPORTING IT, OR EXPORTING IT, & CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE ABOVE PURPOSE, OR USING IT IN LUCING A HYBRID OR DIFFERENT VARIETY THEREFROM, TO THE EXTENT PROVIDED BY THE PLANT VARIETY

CTION ACT. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

CORN, FIELD

'PH4TV'

In Vestimonn Macrest, I have hereunto set my hand and caused the seal of the Plant Invictor Protection Office to be affixed at the City of Washington, D.C. this thirtieth day of January, in the year two thousand two.

Jac M. Jukul

Commissioner Plant Variety Protection Office Amicalysis Washeling Semin l Agriculture

Owner(s) is(are) informed that false representation herein can jeopardize protection and results in penalties.

SIGNATURE OF OWNER

SIGNATURE OF OWNER

NAME (Please print or type)

NAME (Please print or type)

Steven R. Anderson

CAPACITY OR TITLE

DATE

Senior Research
Associate

Associate

INSTRUCTIONS

GENERAL: To be effectively filed with the Plant Variety protection Office (PVPO), ALL of the following items must be received in the PVPO: (1) Completed application form signed by the owner; (2) completed Exhibits A,B,C,E; (3) for a seed reproduced variety at least 2,500 viable untreated seeds, for a hybrid variety sy Irsdy 2,500 untreated seeds of each line necessary to reproduce the variety, or for tuber reproduced varieties ventication that a viable (in the sense that it will reproduce an entire plant) tissue culture will be deposited and maintained in a approved public repository; (4) check drawn on a U.S. bank for \$2,450 (\$300 filing fee and \$2,150 examination fee), payable to "Treasurer of the United States" (See Section 97.6 of the Regulations and Rules of Practice.) Partial applications will be held in the PVPO for not more than 90 days, then returned to the applicant as unfiled. Mail application and other requirements to Plant Variety Protection Office, AMS, USDA, Room 500, NAL Building, 10301 Baltimore Avenue, Beltsville, MD 20705-2351. Retain one copy for your files. All items on the face of the application are self explanatory unless noted below. Corrections on the application form and exhibits must be initialed and dated. DO NOT use masking materials to make corrections. If a certificate is allowed, you will be requested to send a check payable to "Treasurer of the United States" in the amount of \$300 for issuance of the certificate. Certificates will be issued to owner, not licensee or agent.

> Plant Variety Protection Office Telephone: (301)504-5518 FAX: (301)504-5291

Homepage: http://www.ams.usda.gov/science/pvp.htm

ITEM

- 18a. Give: the genealogy, including public and commercial varieties, lines, or clones used, and the breeding method;
 - the details of subsequent stages of selection and multiplication; (2)
 - (3)evidence of uniformity and stability; and
 - (4)the type and frequency of variants during reproduction and multiplication and state how these variants may be identified.
- 18b. Give a summary of the variety's distinctness. Clearly state how this application variety may be distinguished from all other varieties in the same crop. If the new variety is most similar to one variety or a group of related varieties:
 - identify these varieties and state all differences objectively;
 - attach statistical data for characters expressed numerically and demonstrate that these are clear differences; and
 - submit, if helpful, seed and plant specimens of photographs (prints) of seed and plant comparisons which clearly indicate distinctness.
- 18c. Exhibit C forms are available from the PVPO for most crops; specify crop kind. Fill in Exhibit C (Objective Description of Variety) form as completely as possible to describe your variety.
- Optional additional characteristics and/or photographs. Describe any additional characteristics that cannot be accurately conveyed in Exhibit C. Use 18d. comparative varieties as is necessary to reveal more accurately the characteristics that are difficult to describe, such as plant habit, plant disease resistance, etc.
- 18e. Section 52(5) of the Act required applicants to furnish a statement of the basis of the applicant's ownership. An Exhibit E form is available from the PVPO.
- 19. If "Yes" is specified (seed of this variety be sold by variety name only, as a class of certified seed), the applicant may NOT reverse this affirmative decision after the variety has been sold and so labeled, the decision published, or the certificate issued. However, if "No" has been specified, applicant may change the choice. (See Regulations and Rules of Practice, Section 7.103).
- 22. See Sections 41, 42, and 43 of the Act and Section 97.5 of the regulations for eligibility requirements.
- 23. See Section 5.5 of the Act for instructions on claiming the benefit of an earlier filing date.
- 22. CONTINUED FROM FRONT (Please provide the date of first sale, disposition, transfer, or use for each country and the circumstances, if the variety (including any harvested material) or a hybrid produced from this variety has been sold, disposed of, transferred, or used in the U.S. or other countries.)
- 23 CONTINUED FROM FRONT (Please give the country, date of filing or issuance, and assigned reference number, if the variety or any component of the variety is protected by intellectual property right (Plant Breeder's Right or Patent).

Nov. 1, 1999; United States, Canada

NOTES; It is the responsibility of the applicant/owner to keep the PVPO informed of any changes of address or change of ownership or assignment or owner's representative during the life of the application/certificate. There is no charge for filing a change of address. The fee for filing a change of ownership or assignment or any modification of owner's name is specified in Section 97.175 of the regulations. (See Section 101 of the Act, and Sections 97.130, 97.131, 97.175(h) of Regulations and Rules of Practice.)

To avoid conflict with other variety names in use, the applicant should check the variety names proposed by contacting: Seed Branch, AMS, USDA, Room 213, Building 306, Beltsville Agricultural Research Center-East, Beltsville, MD 20705. Telephone: (301) 504-8089.

Public reporting burden for this collection of information is estimated to average 30 minutes per response, including the time for reviewing instruction, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate of any other aspect of this collection of information, including suggestions for reducing this burden, to Department of Agriculture, Clearance Officer, OIRM, AG Box 7630, Jamie L. Whitten Building, Washington, D.C. 20250. When replying, refer to OMB No. 0581-0055 and form number in your letter. Under the PRA of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

The U.S. Department of Agriculture (USDA) prohibits discrimination in its programs on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, and marital or familial status. (Not all prohibited bases apply to all programs). Persons with disabilities who require alternative means for communication of program information (braille, large print, audiotape, etc.) should contact the USDA Office of Communications at (202) 720-2791. To file a complaint, write the Secretary of Agriculture, U.S. Department of Agriculture, Washington, D.C. 20250, or call (202) 720-7327 (voice) or (202) 720-1127 (TDD). USDA is an equal employment opportunity employer.

Exhibit A. Origin and Breeding History

Pedigree: PHKM5<2PHRE1)7322X

200000254

Pioneer Line PH4TV, Zea mays L., a dent-like corn inbred, was developed by Pioneer Hi-Bred International, Inc. from the single cross hybrid PHKM5 (Certificate No. 9400097) X PHRE1 (PVP Certificate No. 9300114) using the backcrossing and pedigree method of plant breeding. Varieties PHKM5 and PHRE1 are proprietary inbred lines of Pioneer Hi-Bred International, Inc. Selfing was practiced (after 2 backcrossing generations) for 5 generations using pedigree selection. During line development, crosses were made to inbred testers for the purpose of estimating the line's combining ability. Yield trials were grown at Moorhead, Minnesota, as well as other Pioneer research locations. After initial testing, additional hybrid combinations have been evaluated and subsequent generations of the line have been grown and hand-pollinated with observations again made for uniformity.

Variety PH4TV has shown uniformity and stability for all traits as described in Exhibit C - "Objective Description of Variety". It has been self-pollinated and ear-rowed 5 generations with careful attention paid to selection criteria and uniformity of plant type to assure genetic homozygousity and phenotypic stability. The line has been increased both by hand and in isolated fields with continued observations for uniformity and stability for 4 generations during the final stages of inbred development and seed multiplication. Very high standards for genetic purity have been established morphologically using field observations and electrophoretically using sound lab molecular marker methodology.

No variant traits have been observed or are expected in PH4TV.

The criteria used in the selection of PH4TV were yield, both per se and in hybrid combinations; late season plant health, grain quality, stalk lodging resistance, and kernel size, especially important in production. Other selection criteria include: ability to germinate in adverse conditions; number of tillers, especially important in production because having numerous tillers increases hybrid production costs spent on detasseling; disease and insect resistance; pollen yield and tassel size.

Exhibit A: Developmental history for PH4TV

	20000254
Season/Year	Inbreeding Level of
Pedigree Grown	Pedigree Grown
Summer 1990	FO
PHKM5, PHRE1	
Winter 1990	F1
PHKM5/PHRE1	
Summer 1992	BC1F1
PHKM5<2PHRE1	
Summer 1993	BC1F2
PHKM5<2PHRE1)7	
Summer 1994	BC1F3
PHKM5<2PHRE1)73	
Summer 1996	BC1F4
PHKM5<2PHRE1)732	
Winter 1996	BC1F5
PHKM5<2PHRE1)7322	
Summer 1997	BC1F6
PHKM5<2PHRE1)73221	
Seed	BC1F6
PHKM5<2PHRE1)7322X	

^{*}PH4TV was selfed and ear-rowed from BC1F2 through BC1F6 generation.
#Uniformity and stability were established from BC1F4 through BC1F6 generation and beyond when seed supplies were increased.

Exhibit B: Novelty Statement

Variety PH4TV mostly resembles Pioneer Hi-Bred International, Inc. proprietary inbred line PHKM5 (PVP Certificate No. 9400097). The data in Tables 1A and 1B are from paired comparisons collected primarily in Johnston and Ankeny, IA. The traits collectively show measurable differences between the two varieties.

Variety PH4TV has wider cob diameter (18.3 mm vs 15.2 mm) than variety PHKM5 (Table 1A, 1B).

Variety PH4TV has less ear weight (27.1 g vs 44.0 g) than variety PHKM5 (Table 1A, 1B).

Variety PH4TV has shorter kernel length (7.1 mm vs 9.2 mm) than variety PHKM5 (Table 1A, 1B).

Variety PH4TV has wider stalk diameter (10.4 mm vs 8.2 mm) than variety PHKM5 (Table 1A, 1B).

Variety PH4TV has longer tassel length (52.3 cm vs 43.9 cm) than variety PHKM5 (Table 1A, 1B).

5/4/21 12/14/0

A t-test was used to compare differences between means and the appropriate parameters have been included. It is difficult to collect standard deviations for table 2 due to the way the historical data was stored.

Exhibit B Novelty Statement Tables

Exhibit B: Novelty Statement Tables

Table 1A: Data from Johnston, IA at 3 different environments in 1999 are supporting evidence for differences between PH4TV and PHKM5 Locations had different environmental conditions. Environments had different planting dates and were in different fields.

Prob (2- tail) Pooled	0.017	0.000	0.000	0.020	0.000	0.000	0.000	0.000	0.002	0.002	0.029	0.005	0.000	0.005	0.002
F-Value Pooled F	2.99	7.20	5.72	-2.88	-5.96	-6.07	-9.19	-5.88	4.43	4.63	2.65	3.89	6.07	3.83	4.53
DF t	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8
StdErro r-2	0.490	0.200	0.316	3.444	1.673	1.435	0.200	0.200	0.245	0.600	0.374	0.510	1.241	1.414	1.806
StdErr or-1	0.548	0.548	0.374	4.261	2.354	2.557	0.200	0.316	0.200	0.245	0.374	0.245	0.927	1.241	0.583
StdDevi	1.095	0.447	0.707	7.701	3.742	3.209	0.447	0.447	0.548	1.342	0.837	1.140	2.775	3.162	4.037
StdDev St	1.225	1.225	0.837	9.529	5.263	5.718	0.447	0.707	0.447	0.548	0.837	0.548	2.074	2.775	1.304
Mean S	2.2	4.2	2.8	-15.8	-17.2	-17.8	-2.6	-2.2	-1.4	3.0	4.1	2.2	9.4	7.2	9.8
Mean- A	15.8	14.8	15.0	47.4	42.0	42.6	9.8	9.2	8.6	8.4	7.8	8.4	43.2	45.0	43.6
	18.0	19.0	17.8	31.6	24.8	24.8	7.2	7.0	7.2	11.4	9.2	10.6	52.6	52.2	52.2
ount- 2	5	ည	2	5	5	5	5	2	5	5	5	5	ಬ	2	5
Count- Count- Mean-	ည	2	5	5	5	2	2	2	2	2	5	2	2	5	5
riety- 2	PHKM5	PHKM5	PHKM5	PHKM5	PHKM5	PHKM5	PHKM5	PHKM5	PHKM5	PHKM5	PHKM5	PHKM5	PHKM5	PHKM5	PHKM5
Variety- Va	PH4TV PH	PH4TV PF	PH4TV PF	PH4TV PF	PH4TV PF	PH4TV PF	PH4TV PF	PH4TV P	PH4TV Ph	PH4TV PH	PH4TV PF	PH4TV PH	PH4TV PF	PH4TV PH	PH4TV PF
Year Traits	1999 cob diameter (mm)	1999 cob diameter (mm)	1999 cob diameter (mm)	1999 ear weight (g)	1999 ear weight (g)	1999 ear weight (g)	1999 kernel length (mm)	1999 kernel length (mm)	1999 kernel length (mm)	1999 stalk diameter (mm)	1999 stalk diameter (mm)	1999 stalk diameter (mm)	1999 tassel length (cm)	1999 tassel length (cm)	1999 tassel length (cm)
Loc Year	20N	岁	Y212	20N	NF	Y212	20N	¥	Y212	20N	볼	Y212	20N	¥	Y212
Station	ΑD	느	ূ	AD	L	ᅩ	AD	느	歬	AD	느	ㅋ	AD	L	H

Table 1B: Summary data from Johnston, IA across environments in 1999 are supporting evidence for differences between PH4TV and PHKM5. Locations had different environmental conditions. Environments had different planting dates and were in different fields. Tables below show means broken out across environments in 1999.

Prob (2- tail) Pooled	0.000	0.000	0.000	0.000	0.000
t-Value Pooled	8.21	-7.14	-9.41	5.47	8.60
DF	28	28	28	28	28
StdErr or-2	0.223	1.418	0.175	0.279	0.831
StdErr or-1	0.300	1.901	0.133	0.289	0.513
	0.862	5.490	0.676	1.082	3.218
StdDevi StdDevi ation-1 ation-2	1.163	7.363	0.516	1.121	1.988
Mean Diff	3.1	-16.9	-2.1	2.2	8.4
Mean- 2	15.2	44.0	9.2	8.2	43.9
Mean-	18.3	27.1	7.1	10.4	52.3
Count- 2	15	15	15	15	15
Count- 1	15	15	15	15	15
Variety- variety-2 Co	V PHKM5	PH4TV PHKM5	PH4TV PHKM5	V PHKM5	PH4TV PHKM5
variet	PH4T	PH4T	PH4T	PH4T	
Lair	1999 cob diameter (mm) PH4TV PHKM5	ear weight (g)	kernel length (mm)	stalk diameter (mm) PH4TV PHKM5	1999 tassel length (cm)
Year	1999	1999	1999	1999	1999

200000254

United States Department of Agriculture, Agricultural Marketing Service Science Division, Plant Variety Protection Office National Agricultural Library Building, Room 500 Beltsville, MD 20705

Objective Description of Variety Corn (Zea mays L.)

Name of Ap	plicant (s)		Variety Seed Source	Variet	y Name or Temporary Designation			
Pioneer I	Hi-Bred Inte	ernational, Inc.	•		PH4TV			
A 11 (G)		TD 11 G': G: T' G 1 1		TOD OFFICIAL TION	·			
		FD No., City, State, Zip Code and	Country	FOR OFFICIAL USE	_			
7301 NW	62 nd Avenu	ie, P.O. Box 85,		PVP0 Number				
Johnston	, Iowa 5013	31-0085		PVPU Number				
Place the appropriate number that describes the varietal characters typical of this inbred variety in the spaces below. Right justify whole numbers by adding Leading zeroes if necessary. Completeness should be striven for to establish an adequate variety description. Traits designated by an '*' are considered Necessary for an adequate variety description and must be completed. COLOR CHOICES (Use in conjunction with Munsell color code to describe all color choices: describe #25 and #26 in Comments section):								
01=Light Gr		06=Pale Yellow	11=Pink	16=Pale Purple	21=Buff			
02=Medium		07=Yellow	12=Light Red	17=Purple	22=Tan			
03=Dark Gre	een	08=Yellow Orange	13=Cherry Red	18=Colorless	23=Brown			
04=Very Dan	rk Green	09=Salmon	14=Red	19=White	24=Bronze			
05=Green-Y	ellow	10=Pink-Orange	15=Red & White	20=White Capped	25=Variegated (Describe) 26=Other (Describe)			
STANDARD	INBRED CHO	DICES						
(Use the mos	st similar (in bac	ckground and maturity) of these to	make comparisons based on g	row-out trial data):				
Yellow Dent	Families:		Yellow Dent (Unrelated):	Sweet Co	orn:			
Family	Members		Co109, ND246,	C13, Io	wa5125, P39, 2132			
B14	CM105, A632,	B64, B68	Oh7, T232,					
B 37	B37, B76, H84		W117, W153R,	Popcorn:				
B73	N192, A679, B	73, NC268	W18BN	SG1533	3, 4722, HP301, HP7211			
	Mo17, Va102,							
Oh43	A619, MS71, H	199, Va26	White Dent:	Pipecorn	:			
WF9	W64A, A554, A	A654, Pa91	C166, H105, Ky228	Mo15W	, Mo16W, Mo24W			

7F9 W64A, A554, A654, Pa91 Groups on Lynx/Osborn/Grunst/98-99PVP



1. TYPE:	(describe intermediate types in Comme	nts section):			Stand	ard Variety	/ Name
2		A554					
2. REGIO	ON WHERE DEVELOPED IN THE U.S.A	A.:			Stand	ard Seed	Source
2	1=Northwest 2=Northcentral 3=Northe	ast 4=Southeast 5=Sou	thcentral			AMES 19	305
	6=Southwest 7=Other					ANNEO 10	500
3. MATU	RITY (In Region of Best Adaptability; sh	ow Heat Unit formula in '	Comments' se	ection)			
DAYS	HEAT UNITS				DAYS	HEAT UN	ITS
065	1,172.3 From emergence to 50% of	plants in silk			065	1,159.3	
065	1,164.3 From emergence to 50% of	plants in pollen			066	1,169.0	
003	0,077.0 From 10% to 90% pollen st	ned			003	0,071.0	
	From 50% silk to optimum e	edible quality					
	From 50% silk to harvest at	25% moisture					
4. PLAN	:		Standard	Sample		Standard	Sampl
			Deviation	Size		Deviation	Size
162.5	cm Plant Height (to tassel tip)		12.02	02	163.0	05.29	03
063.5	cm Ear Height (to base of top ear nod	e)	09.19	02	055.7	05.77	03
012.3	cm Length of Top Ear Internode		01.89	03	011.4	00.53	03
0.0	Average Number of Tillers/plant		00.05	03	0.0	00.03	03
0.4	Average Number of Ears per Stalk		00.31	03	1.0	00.11	03
3	Anthocyanin of Brace Roots: 1=Abse	nt 2=Faint 3=Moderate	4=Dark		3		
5. LEAF:			Standard	Sample		Standard	Sample
			Deviation	Size		Deviation	Size
07.6	cm Width of Ear Node Leaf		00.35	03	09.1	00.31	03
72.3	cm Length of Ear Node Leaf		01.17	03	66.9	04.11	03
04			00.35	03	05	00.60	03
17	Degrees Leaf Angle (measure from 2n at anthesis to stalk above leaf)	d leaf above ear	00.42	03	22	04.41	03
03	Leaf Color (Munsell code)	5GY34			03	5G`	/44
1	Leaf Sheath Pubescence (Rate on sca	le from 1=none to 9=like	peach fuzz)		1		
	Marginal Waves (Rate on scale from 1:	none to 9=many)					
	Longitudinal Creases (Rate on scale from	om 1=none to 9=many)					
6. TASSE	L:		Standard	Sample		Standard	Sample
	· 		Deviation	Size		Deviation	Size
05	Number of Primary Lateral Branches		00.83	03	12	01.59	03
26	Branch Angle from Central Spike		00.81	03	19	05.35	03
52.3	cm Tassel Length (from top leaf collar	to tassel tip)	00.23	03	45.9	02.39	03
	Pollen Shed (rate on scale from 0=mal		l)		5		
14	Anther Color (Munsell code)	2.5R36			07	5Y	'88
01	Glume Color (Munsell code)	5GY48			01	5G	Y66
1	Bar Glumes (Glume Bands): 1=Absent	2=Present			1		
Annlineti-	n Variaty Data	Page 1			Standa	rd Variety	Data
∠hhiicano	n Variety Data	raye i			Sidilud	ia vallety	Data

pplication	Variety Data PH4TV	Page 2			Standa	rd Variet	y Data	
7a. EAR ((Unhusked Data):							
<u>14</u>	Silk Color (3 days after emergence) (N	/lunsell code)		10RP28	<u>07</u>	<u>2.5G\</u>	<u> 196</u>	
<u>01</u>	Fresh Husk Color (25 days after 50% s	silking) (Munsell code)		<u>5GY68</u>	<u>01</u>	5GY	<u>78</u>	
<u>21</u>	Dry Husk Color (65 days after 50% silk	Ory Husk Color (65 days after 50% silking) (Munsell code) 2.5Y92						
<u>3</u>	Position of Ear at Dry Husk Stage: 1= 0	Upright 2= Horizontal	3= Pendant		<u>3</u>			
1	Husk Tightness (Rate of Scale from 1=	very loose to 9=very t	ight)		<u>7</u>			
<u>2</u>	Husk Extension (at harvest): 1=Short (ears exposed) 2=Med	ium (<8 cm)		<u>2</u>			
	3=Long (8-10 cm beyond ear tip) 4=Ve	ery Long (>10 cm)						
7b. EAR	(Husked Ear Data):		Standard	Sample	Star	ndard	Sampl	
			Deviation	Size	Devi	iation	Size	
<u>11.0</u>	cm Ear Length		00.00	<u>03</u>	<u>09.7</u> <u>0</u>	00.58	<u>03</u>	
<u>29.3</u>	mm Ear Diameter at mid-point		<u>00.58</u>	<u>03</u>	39.3 C)1. <u>15</u>	<u>03</u>	
027.3	gm Ear Weight		<u>04.04</u>	<u>03</u>	67.0 <u>0</u>	6.24	<u>03</u>	
<u>15</u>	Number of Kernel Rows		<u>01.15</u>	<u>03</u>	<u>14.0</u> <u>0</u>	00.00	<u>03</u>	
<u>2</u>	Kernel Rows: 1=Indistinct 2=Distinct				<u>2</u>			
1	Row Alignment: 1=Straight 2=Slightly 0	Curved 3=Spiral			1			
<u>17.3</u>	cm Shank Length		<u>01.15</u>	<u>03</u>	<u>11.7</u> <u>0</u>	<u>)3.79</u>	<u>03</u>	
2	Ear Taper: 1=Slight 2= Average 3=Extr	reme			2			
8. KERNE	EL (Dried)		Standard	Sample	Standa	ard	Sampl	
			Deviation	Size	Deviati	ion	Size	
<u>07.0</u>	mm Kernel Length		00.00	<u>03</u>	<u>10.3</u> 0	0.58	<u>03</u>	
<u>07.0</u>	mm Kernel Width		00.00	<u>03</u>	08.0 0	0.00	<u>03</u>	
04.0	mm Kernel Thickness		00.00	<u>03</u>	<u>04.0</u> 0	0.00	<u>03</u>	
00.0	% Round Kernels (Shape Grade)		00.00	<u>03</u>	<u>28.7</u> 0	<u>2.31</u>	<u>03</u>	
1	Aleurone Color Pattern: 1-Homozygous	s 2=Segregating			1			
<u>07</u>	Aluerone Color (Munsell code)		<u>10'</u>	YR714	<u>07</u>	2.5Y	<u> 312</u>	
<u>07</u>	Hard Endosperm Color (Munsell code)		<u>10</u>	YR712	<u>07</u>	2.5Y	<u> 312</u>	
<u>03</u>	Endosperm Type:				<u>3</u>			
	1=Sweet (Su1) 2=Extra Sweet (sh2) 4=High Amylose Starch 5=Waxy Sta 7=High Lysine 8=Super Sweet (se) 10=Other	arch 6=High Protein						
<u>09.0</u>	gm Weight per 100 Kernels (unsized sa	mple)	01.00	<u>03</u>	<u>22.00</u> 0	2.65	<u>03</u>	
9. COB:	<u> </u>		Standard	Sample	St	andard	Samp	
			Deviation	Size	}	eviation	Size	
<u>18</u> .3	mm Cob Diameter at mid-point		00.58	<u>03</u>	<u>24.3</u>	<u>00.58</u>	<u>03</u>	
	Cob Color (Munsell code)	<u>5Y91</u>		_	14	 10F		

Application Variety Data

Page 3

Standard Variety Data

40 DIOEAGE DI	CONTANIOS (Data fivere 4 (secret propositive) to 0 (secret registered).		
	ESISTANCE (Rate from 1 (most susceptible) to 9 (most resistant); if not tested; leave Race or Strain Options blank if polygenic):		
A. Leaf Bl	ights, Wilts, and Local Infection Diseases		
	Anthracnose Leaf Blight (Colletotrichum graminicola)		
	Common Rust (Puccinia sorghi)		
	Common Smut (Ustilago maydis)		
<u>6</u>	Eyespot (Kabatiella zeae)	3	
<u>4</u>	Goss's Wilt (Clavibacter michiganense spp. nebraskense)	Z	
	Gray Leaf Spot (Cercospora zeae-maydis)		
	Helminthosporium Leaf Spot (Bipolaris zeicola) Race ——		
	Northern Leaf Blight (Exserohilum turcicum) Race ——		
	Southern Leaf Blight (Bipolaris maydis) Race ——		
	Southern Rust (Puccinia polysora)		
<u>2</u>	Stewart's Wilt (Erwinia stewartii)	4	
	Other (Specify) ———		
B. System	nic Diseases		
	Corn Lethal Necrosis (MCMV and MDMV)		
<u>4</u>	Head Smut (Sphacelotheca reiliana)	4	
	Maize Chlorotic Dwarf Virus (MDV)		
	Maize Chlorotic Mottle Virus (MCMV)		
	Maize Dwarf Mosaic Virus (MDMV)		
	Sorghum Downy Mildew of Corn (Peronosclerospora sorghi)		
	Other (Specify) ———		
C. Stalk F	tots		
	Anthracnose Stalk Rot (Colletotrichum graminicola)		
	Diplodia Stalk Rot (Stenocarpella maydis)		
	Fusarium Stalk Rot (Fusarium moniliforme)		
	Gibberella Stalk Rot (Gibberella zeae)		
	Other (Specify) ——		•
D. Ear an	d Kernel Rots		
	Appercially Ear and Vernel Pet (Appercially flagge)		
	Aspergillus Ear and Kernel Rot (Aspergillus flavus) Diplodia Ear Rot (Stenocarpella maydis)		
	Fusarium Ear and Kernel Rot (Fusarium moniliforme)	_	
7	Gibberella Ear Rot (Gibberella zeae)	<u>6</u>	
<u>7</u>	Other (Specify) ———	<u> </u>	
	Outer (Openity)		

Application Variety Data

Page 4

Standard Variety Data

	tate how heat units were Continue in Exhibit D):	e calculated, standard inbred se	ed source, a	and/or where	
	1 Isozymes	Q RFLP's	<u>0</u>		
13. MOLECU	LAR MARKERS: (0=dat	a unavailable; 1=data available	but not sup	plied; 2=data supplied)	:
<u>2,480.0</u>	Kg/ha Yield of Inbred P	er Se (at 12-13% grain moisture	;)	<u>2,259.3</u>	
<u>4.0</u>		Iging (at 65 days after anthesis)		38.3	
	% Pre-anthesis Root Lo	~ ~			
	% Pre-anthesis Brittle S	•			
<u>0.0</u>	% Dropped Ears (at 65	•		0.0	
_	on a scale from 1=wors	st to excellent)			
12. AGRON	Staygreen (at 65 days	after anthesis) (Rate		2	
12 AGRON	OMIC TRAITS:				
	Other (Specify) ———				
		abrotica virgifrea virgifera)			
		te (Tetranychus urticae)			
	cm tunneled/plant				
	Stalk Tunneling				
	Leaf Feeding				
	Southwestern Corn Bo	rer (Diatreaea grandiosella)			
	Southern Rootworm (D	iabrotica undecimpunctata)			
	Northern Rootworm (D	iabrotica barberi)			
	Maize Weevil (Sitophile	us zeamaize			
	mg larval wt.				
	Silk Feeding				
	Leaf Feeding				
	Fall Armyworm (Spodo	ptera fruqiperda)		·	
	cm tunneled/plant				
	Stalk Tunneling				
	2nd Generation (Typ	ically Leaf Sheath-Collar Feedir	ng)		
	•	cally Whorl Leaf Feeding)			
	European Corn Borer (•			
	Corn Sap Beetle (Carp	•			
	Corn Leaf Aphid (Rhor	palosiphum maidis)			
	Ear Damage				
	mg larval wt.				
	Silk Feeding				
	Corn Worm (Helicover Leaf Feeding	pa zea)			
	Corn Morm (Halicovar	na 70a)			

Application Variety Data

Page 4

Standard Variety Data

CLARIFICATION OF DATA IN EXHIBITS B AND C

Please note the data presented in Exhibit C, "Objective Description of Variety," are collected primarily at Johnston and Ankeny, Iowa. The data in Exhibit B are from comparisons of inbreds grown in the same tests in Johnston and Ankeny, IA. The data in Tables 1A and 1B are from paired comparisons collected in Johnston and Ankeny, IA. These traits collectively show distinct differences between the two varieties.

The data collected in exhibit C were collected from environments 1999 for page 1 and 2. There are factors that differ from environment to environment. The environments had different planting dates. Environmental temperature and precipitation differences during the vegetative and grain fill periods can impact plant and grain traits and be a source of variability. These data are mostly based on 5 plants measured at each location. There often is more variability associated with environment to environment factors than within locations. Please see Table 3 for average temperature and rainfall information in 1999. In addition, this variety showed susceptibility to Stewarts Wilt in 1999. A high population of flea beetles in 1999 at 2 of our locations may have contributed more variability than usual resulting in higher standard deviations for some traits.

The number of ears per stalk was small probably due to susceptibility to Stewart's Wilt disease that caused many of the plants to go barren. One location in particular had high flea beetle populations and severe Stewart's Wilt. Similar flea beetle populations in 2000 also resulted in a similar low value for ears per stalk.

Notes from our field book indicate that Stewart's Wilt may have caused poorer seed quality than normal in at least 1 of our locations. Poor seed quality probably resulted in lower kwt/100k than normal. Although we also had high levels of flea beetle pressure in 2000 the average kwt/100k in that year was 14 g/100k.

Table 3. Temperature and Rainfall

TEMPERATURE

YEAR	MAY	JUN	JULY	AUG	AVERAGE
1994	59.8	70.7	71.9	69.0	67.9
1995	56.2	69.4	74.3	76.9	69.2
1996	56.2	69.3	71.3	70.5	66.8
1997	53.5	70.6	74.1	69.6	67.0
1998	64.7	66.6	74.8	73.5	69.9
1999	60.7	69.7	78.7	70.5	69.9

RAINFALL

YEAR	MAY	JUN	JULY	AUG	Total
1994	3.67	5.75	1.71	4.18	15.31
1995	5.04	4.19	2.94	2.87	15.04
1996	8.47	4.35	2.51	2.14	17.47
1997	4.32	3.27	4.10	1.36	13.05
1998	6.46	11.07	5.70	4.96	28.19
1999	6.46	4.54	4.45	6.55	21.85

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE	The following statements are made in acco	
EXHIBIT E STATEMENT OF THE BASIS OF OWNERSHIP	Application is required in order to determ certificate is to be issued (7 U.S.C. 2421). until certificate is issued (7 U.S.C. 2426).	nine if a plant variety protection Information is held confidential
1. NAME OF APPLICANT(S)	2. TEMPORARY DESIGNATION	3. VARIETY NAME
PIONEER HI-BRED INTERNATIONAL, INC.	OR EXPERIMENTAL NUMBER	PH4TV
4 .ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP, and Country)	5. TELEPHONE (include area code)	6. FAX (include area code)
7301 NW 62 nd AVENUE	515-270-4051	515-253-2125
P.O.BOX 85	7. PVPO NUMBER 0 0 0 0 2) F ₁ /.
JOHNSTON, IA 50131-0085		. O 4
8. Does the applicant own all rights to the variety? Mark an "X" in appropriate bl	ock. If no, please explain: 🛛 YES	□NO
9. Is the applicant (individual or company) a U.S. national or U.S. based compar	ny? ☑ YES ☐ NO	
If no, give name of country		
10. Is the applicant the original owner?	please answer <u>one</u> of the following:	
 a. If original rights to variety were owned by individual(s), is(are) the orig 	inal owner(s) a U.S. national(s)?	
☐ YES ☐ NO if no, give name of country		
b. If original rights to variety were owned by a company(ies), is(are) the	original owner(s) a U.S. based company?	
☑ YES ☐ NO If no, give name of country		
11. Additional explanation on ownership (if needed, use reverse for extra space):		
PH4TV is owned by Pioneer Hi-Bred International, Inc.		
PLEASE NOTE:		
Plant variety protection can be afforded only to owners (not licensees) who meet one of the	ne following criteria:	
 If the rights to the variety are owned by the original breeder, that person must be a Which affords similar protection to nationals of the U.S. for the same genus and specific 		ntry, or national of a country
2. If the rights to the variety are owned by the company which employed the original becountry, or owned by national of a country which affords similar protection to nation		
3. If the applicant is an owner who is not the original owner, both the original owner a	and the applicant must meet one of the above cri	teria.
The original breeder/owner may be the individual or company who directed final breeding	g. See section 41(a)(2) of the Plant Variety Pro-	tection Act for definition.

According to the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this information collection is 0581-0055. The time required to compete this information collection is estimated to average 10 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in its programs on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, and marital or familial status (Not all prohibited bases apply to all programs). Persons with disabilities who require alternative means for communication of program information (braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

Center at 202-720-2600 (voice and TDD).

To file a complaint, write Secretary of Agriculture, U.S. Department of Agriculture, Washington, D.C. 20250, or call 1-800-245-6340 (voice) or (202) 720-1127 (TDD) USDA is an equal employment